

Literature

The reference list is in common for all three data bases, although it appears as part of each data base. This is simply for practical purposes.

LITERATURE REFERENCES TO DELILA I, DELILA II and DELILA III

[1] Albrektson A. Needle litterfall in stands of *Pinus sylvestris* L. in Sweden, in relation to site quality, stand age, and latitude. Scand. J. For. Res. 3 (1988) 333-342.

[2] Anonymous. Redovisning av fasta försöksytor. Royal College of Forestry, Department of Forest Yield Research, Research Notes, No 33 Stockholm, (1974). (In Swedish).

[3] Aussenac G. Production de litiere dans divers peuplements forestiere de l'est de la France. Oecologia Plantarum 4 (1969) 225-236.

[4] Berg B., Johansson M.-B., and Lundmark J.-E. Site descriptions for experimental forest sites - a compilation. Department of Forest Ecology and Forest Soils. Swedish University of Agri-cultural Sciences, Uppsala. Research Report No 72, 1997, 43 pp. ISRN SLU-SKOMA-R--73--SE.

[5] Berg B., Berg M., Flower-Ellis J.G.K., Gallardo A., Johansson M., Lundmark J.-E., and Madeira M. Amounts of litterfall in some European Coniferous Forests, in: Brey-meyer, A. (Ed.). Proceedings from Scope Seminar. Conference Papers 18. Geogra-phy of Carbon Budget Processes in Terrestrial Ecosystems. Szymbark, Aug. 17-23, 1991,(1993) pp 123-146.

[6] Berg, B., Albrektson, A., Berg, M., Cortina, J., Johansson, M.-B., Gallardo, A., Madeira, M., Pausas, J., Kratz, W., Vallejo, R., and McClaugherty, C. 1999. Amounts of litter-fall in pine forests in the northern hemisphere, especially Scots pine. Annals of Forest Science 56;625-639.

[7] Berg, B., Johansson, M., Tjarve, I., Gaitnieks, T., Rokjanis, B., Beier, C., Rothe,

A., Bolger, T., Göttlein, A., and Gerstberger, P. 1999. Needle litterfall in a north European spruce forest transect. Reports from the Departments of Forest Ecology and Forest Soils, Swedish University of Agricultural Sciences. Report 80, 54 pp.

[8] Bonnevie-Svendsen C., and Gjems O. Amount and chemical composition of the litter from larch, beech, Norway spruce and Scots pine stands and its effect on the soil. *Meddelelser fra det Norske skogsforsöksvesen* 14,(48)(1957) 115-168.

[9] Zimmermann, S., Braun, S., Conedera, M., and Blaser, P. 2002. Macronutrient inputs by litter fall as opposed to atmospheric deposition into two contrasting chestnut forest stands in southern Switzerland. *Forest Ecology and Management* 161; 289-302.

[10] Casals P., Romanyo J., Cortina J., Fons J., Bode M., and Vallejo V.R. Nitrogen supply rate in Scots pine (*Pinus sylvestris* L.) forests of contrasting slope aspect. *Plant and Soil* 168/169, (1995) 67-73.

[11] Persson, T., and Nilsson, L.O. (Eds) Skogabyförsöket - effekter av långvarig kväve- och svaveltillförsel till ett skogssystem. Naturvårdsverket Rapport 5173. Naturvårdsverkets förlag. Stockholm. ISBN 90-620-5173-3. (In Swedish).

[12] Falck J. Changes in the nutrient content of vegetation and forest floor after fertilization with urea in a mature Scots pine stand (*Pinus sylvestris* L.). Swedish University of Agricultural Sciences, Department of Silviculture, Report 5 (1981).

[13] Flower-Ellis J.G.K. Litterfall in an age series of Scots pine stands: Summary of results for the period 1973-1983. Department of Ecology and Environmental Research. Swedish University of Agricultural Sciences. Report 19 (1985) 75-94. ISSN 0348-422X.

[14] Flower-Ellis J.G.K., and Olsson L. Litterfall in an age series of Scots pine stands and its variation by components during the years 1973-1978. Swedish University of Agricultural Sciences. Swedish Coniferous Forest Project. Technical Report No 15 (1978). ISSN 0346-7708

[15] Gloaguen J.C., and Touffet J. Production de litiere et apport au sol delements mineraux dans quelques peuplements resineaux de Bretagne. *Annal.Sci. Forest.* 33 (1976) 87-107. (In French, English summary).

[16] Hanssen, K., and Bastrup-Birk, A. (Eds) 200X. Naeringsstofkredsløb i skove - ionbalanceprojektet. Chapter 4. Biomasse, produktion og CO₂-optagelse. (In Danish).

[17] Helmisaari H.-S. Nutrient cycling in *Pinus sylvestris* stands in eastern Finland. *Plant and Soil* 168-169 (1995) 327-336.

[18] Nihlgård B. 1972. Plant biomass, primary production and distribution of chemical elements in a beech and a planted spruce forest in South Sweden. *Oikos* 23; 69-81.

[19] Mälkönen E. Annual primary production and nutrient cycle in some Scots pine stands. *Commun. Inst. Forest. Fenn.* 84,5 (1974).

[20] Mälkönen E., and Kukkola M. Effect of long-term fertilization on the biomass production and nutrient status of Scots pine stands. *Fertilizer Research* 27 (1991) 113-127.

[22] Pausas J.G. Litter fall and litter decomposition in *Pinus sylvestris* forests of the eastern Pyrenees. *J. Vegetat. Sci.* 8 (1997) 643-650

[23] Rodin L.E., and Basilewicz N.I. Production and mineral cycling in terrestrial vegetation. Oliver and Boyd, London, 1967.

[24] Solberg, S., Berg, I.A., Horntvedt, R., Aamlid, D., and Torseth, K. 1997. Intensive Skogsovervåkningsoverflater. Resultater fra 1995. (Intensive forest monitoring plots. Results 1995). *Aktuellt fra Skogforsk.* 4

[25] Faituri MY. 2001. Soil Organic Matter in Mediterranean and Scandinavian forest ecosystems and dynamics of nutrients and monomeric phenolic compounds.

Silvestra 236, 136 pp.

[26] Solberg, S., Berg, I.A., Brevik K., Groeggen, T., Moshaug, E., Torseth, K., and, Aamlid, D. 1998. Intensive Skogsovervåkningsoverflater. Resultater fra 1997. (Intensive forest monitoring plots. Results 1997). Aktuellt fra Skogforsk. 5

[27] Santa Regina, I., Gallardo, J.F. 1985. Produccion de hojaresca en tres bosques de la sierra de Bejar (Salamanca). Mediterranea Ser. Biol. 8; 89-101 (In Spanish).

[28] IFEF. The IFEF data base. provisional reference.

[29] Matzner, E., 1988. Ser Stoffumsatz zweier Waldökosysteme im Solling. Berichte des Forschungszentrums Waldökosysteme/Waldsterben. A40:1-217. (In German).

[30] Matzner, E., Khanna PK, Meiwes, KJ, Lindheim, J Prenzel, J., Ulrich, B. 1982. Elementfluesse in Waldökosystemen im Solling - Datendokumentation. Göttinger Bodenkundliche Berichte 71: 1-267. (In German).

[31] Kreutzer, K., Göttlein, A. 1991. Ökosystemforschung Höglwald. Beiträge zur Auswirkung von saurer Beregnung und Kalkung in einem Fichtenaltbestand. Heft 39. (In German).

[32] Kreutzer, K., Bittersohl, J. 1986. Anwirkungen des sauren Regens und der kompensatorischen Kalkung im Höglwald. Forstwissenschaftl. Centralbl. 105: 273-282. (In German).

[33] Berg, B., Gerstberger, P. 2004. Element fluxes with litter fall in mature old stands of Norway spruce and European beech in Bavaria, south Germany. In; Matzner, E, (Ed) Biogeochemistry of Forested Catchments in a Changing Environment. A German Case Study. Ecological Studies Vol 172 pp 271-278. Springer Verlag, Berlin - Heidelberg

[34] Malek, S. 2002. Litterfall in beech stands as a source for elements - an example of a 5 years study on monitoring plost in Ojcow National Park and Forest Experimental

Station in Krynica - Southern Poland. pp 114-123. In: Ed L. Björk. Sustainable Forestry in temperate regions. Proceedings of the SUFOR International Workshop, April 7-9, 2002 in Lund Sweden. Reports in Ecology and Environmental Engineering. No 1.

[35] Newbould P.J. 1967. Methods for estimating the primary production of forests IBP Handbook No 2. Blackwell, Oxford.

[36] Flower-Ellis JGK. 1985. Litterfall in an age series of Scots pine stands: Summary of results from the period 1973-1983, Department of Ecology and Environmental Research. Swedish University of Agricultural Sciences. Report 19, pp 75-94.

[40] Howard, P. J. A., and Howard, D. M. 1974. Microbial decomposition of tree and shrub leaf litter. 1. Weight loss and chemical composition of decomposing litter. *Oikos* 25:311-352.

[41] Berg, B. 1998. A maximum limit for foliar litter decomposition - a synthesis of data from forest systems. Reports from the Departments of Forest Ecology and Forest Soils, Swedish University of Agricultural Sciences. Report 77, 158 pp.

[42] Berg, B., Ekbohm, G. 1991. Litter mass-loss rates and decomposition patterns in some needle and leaf litter types. Long-term decomposition in a Scots pine forest VII. *Canadian Journal of Botany* 69:1449-1456.

[43] Berg, B., Ekbohm, G. 1993. Decomposing needle litter in Lodgepole pine (*Pinus contorta*) and Scots pine (*Pinus sylvestris*) monocultural systems. Is there a maximum mass loss? *Scandinavian Journal of Forest Research*. 8:457-465.

[44] Berg, B. 2000. Initial rates and limit values for decomposition of Scots pine and Norway spruce needle litter - a synthesis for N-fertilized forest stands. *Canadian Journal of Forest Research* 30; 122 -135.

[45] Berg, B., Ekbohm, G., Johansson, M.-B., McClaugherty, C., Rutigliano, F., Virzo De Santo, A. 1996. Maximum decomposition limits of forest litter types; a synthesis. *Canadian Journal of Botany* 74:659-672.

[46] Berg, B., Virzo De Santo, A., Rutigliano, F., and Ekbohm, G. 2003. Limit values for plant litter decomposing in two contrasting soils - influence of litter elemental composition. *Acta Oecologica* 24:295-302.

[47] Davey, M., Berg, B., Emmett, B., and Rowland, P. 200X. Decomposition of oak leaf litter is related to initial litter Mn concentrations. *Canadian Journal of Botany* XX; XXX-XXX.

[48] De Angelis, P., Chigwerewe, K.S., and Scarasia Mugnozza, G.E. 2000. Litter quality and decomposition in a CO₂-enriched Mediterranean forest ecosystem. *Plant and Soil* 224: 31-41.

[49] Prof Francesca Cotrufo, personal communication

[50] Starr, M., Saarsalmi, A., Hokkanen, T., Merilä, P., Helimsaari, H-S. 2005. Models of Litterfall production for Scots pine (*Pinus sylvestris* L.) in Finland using stand, site and climate factors. *Forest Ecology and Management* 205: 215-225.

[51] Saarsalmi, A., Starr, M., Hokkanen, T., Ukonmaanaho, L., Kukkola, M., Nöjd, P., Sievänen, R. 2007. Predicting annual canopy litterfall production for Norway spruce (*Picea abies* (L) Karst.) stands. *Forest Ecology and Management* 242: 578-586.